


CINNAMIC ACID
MONOPHENOL MONOOXYGENASE

KK.
FF 78/04.
Sug
p

SKRIPSI

TITA SUGESTI

PERBANDINGAN AKTIVITAS ASAM SINAMAT, ASAM 3-HIDROKSISINAMAT, DAN ASAM 4-HIDROKSISINAMAT SEBAGAI TIROSINASE INHIBITOR



**FAKULTAS FARMASI UNIVERSITAS AIRLANGGA
BAGIAN KIMIA FARMASI
SURABAYA
2003**

Lembar Pengesahan

**PERBANDINGAN AKTIVITAS ASAM SINAMAT,
ASAM 3-HIDROKSISINAMAT, DAN
ASAM 4-HIDROKSISINAMAT
SEBAGAI TIROSINASE INHIBITOR**

SKRIPSI

**Dibuat Untuk Memenuhi Syarat
Mencapai Gelar Sarjana Farmasi Pada Fakultas Farmasi
Universitas Airlangga
2003**

Oleh :

**Tita Sugesti
NIM : 059912171**

**Skripsi ini telah disetujui
tanggal 28 Oktober 2003 oleh :**

Pembimbing Utama



Dr. Tutuk Budlati, MS.
NIP. 130 531 780

Pembimbing Serta



Hadi Poerwono, MSc., PhD.
NIP. 131 831 452

ABSTRACT

The investigation of inhibition of tyrosinase activity by cinnamic acid, 3-hydroxycinnamic acid, and 4-hydroxycinnamic acid was done using L-Tyrosine as substrate.

The percentage of tyrosinase inhibition activity was measured using UV-Vis spectrophotometer that give the absorbance of dopachrome produced by the reaction of L-Tyrosin, tyrosinase, and the inhibitor.

Michaelis Menten constant (K_M) and maximal velocity (V_{max}) of the tyrosinase was determined by Lineweaver-Burk's plots. The value of K_M and V_{max} which obtained were 0,961 mM and 0,020 $\mu\text{mol/min}$. Lineweaver-Burk's plot of several concentration of L-Tyrosine showed that cinnamic acid inhibited tyrosinase activity non competitively but the 3-hydroxycinnamic acid and 4-hydroxycinnamic acid inhibited competitively.

The percentage of tyrosinase activity of cinnamic acid, 3-hydroxycinnamic acid, and 4-hydroxycinnamic acid were 14,0%, 12,6%, and 79,1% respectively.

Key word : cinnamic acid, 3-hydroxycinnamic acid, 4-hydroxycinnamic acid, tyrosinase, inhibition activity.